

Antimicrobial resistance among human non-typhoidal *Salmonella*: A comparison between Enter-net in Europe and NARMS in the United States

J.E. Stevenson¹, E.J. Threlfall², T.J. Barrett¹, T.M. Chiller¹

1. National Center for Infectious Disease, Centers for Disease Control and Prevention, Division of Bacterial and Mycotic Diseases, Foodborne and Diarrheal Diseases Branch, NARMS Unit
2. Health Protection Agency, Specialist and Reference Microbiology Division, Laboratory of Enteric Pathogens, London, United-Kingdom (on behalf of Enter-net participating countries)

Background: Antimicrobial resistance, including multidrug resistance, contributes to the human health burden of *Salmonella* and may compromise the efficacy of available treatment options. In past decades, multidrug-resistant phenotypes of non-typhoidal *Salmonella* have disseminated in Europe and the United States. Efforts to mitigate dissemination of multidrug-resistant strains are enhanced by international collaborations.

Methods: In 2000, antimicrobial susceptibility results for human non-typhoidal *Salmonella* isolates were submitted from 10 European countries (pop. 280 million) to Enter-net. In 2000, isolates of human non-typhoidal *Salmonella* were submitted from 15 states in the United States (pop. 108 million) to the National Antimicrobial Resistance Monitoring System (NARMS). Results for Enter-net and NARMS were compared.

Results: Enter-net received results from 27,059 isolates of human non-typhoidal *Salmonella*; NARMS received and tested 1,367 human non-typhoidal *Salmonella* isolates. Higher proportions of sulfonamide (30% vs. 17%), tetracycline (26% vs. 19%), and nalidixic acid (14% vs. 2%) resistance were observed in Enter-net than in NARMS; a higher proportion of NARMS isolates (3%) were resistant to a 3rd-generation cephalosporin (ceftiofur) than Enter-net isolates (0.6% - cefotaxime). The proportion of isolates that were fluoroquinolone-resistant was similar in both systems, 0.5% in Enter-net and 0.4% in NARMS. *S. Typhimurium* comprised 25% of the Enter-net and 22% of the NARMS collection. A higher proportion of *S. Typhimurium* isolates were R-type ACSSuT in Enter-net (50%) than in NARMS (28%). *S. Newport* comprised 1% of the Enter-net and 9% of the NARMS collection. A higher proportion of *S. Newport* isolates were R-type ACSSuT in NARMS (23%) than Enter-net (5%).

Conclusion: Resistance to clinically important antimicrobials among human non-typhoidal *Salmonella* is common in both Europe and the United States. Substantial differences exist in the patterns of specific resistances between the two regions. Understanding the emergence and dissemination of antimicrobial-resistant *Salmonella* is strengthened by international collaborations.